



# Near-Earth plasma sheet as a seed population for the outer radiation belt

**Natalia Ganushkina (1, 2) and Stepan Dubyagin (1)**

*(1) Finnish Meteorological Institute, Helsinki, Finland*

*(2) University of Michigan, Ann Arbor MI, USA*

***Special thanks to Dr. Elena Grigorenko (Space Research Institute, Moscow, Russia)  
for Cluster data analysis***

The research leading to these results was partly funded by the European Union Seventh Framework Programme (FP7/2007-2013) under grant agreement No 606716 SPACESTORM

Cluster 15th and Double Star 10th Anniversary  
Workshop 12---16 October 2015 Venice, Italy



**British  
Antarctic Survey**  
NATURAL ENVIRONMENT RESEARCH COUNCIL

**ONERA**  
THE FRENCH AEROSPACE LAB



ILMATIETEEN LAITOS  
METEOROLOGISKA INSTITUTET  
FINNISH METEOROLOGICAL INSTITUTE





# Near-Earth plasma sheet as a seed population for the outer radiation belt

Natalia Ganushkina (1, 2) and Stepan Dubyagin (1)

(1) *Finnish Meteorological Institute, Helsinki, Finland*

(2) *University of Michigan, Ann Arbor MI, USA*

The research leading to these results was partly funded by the European Union Seventh Framework Programme (FP7/2007-2013) under grant agreement No 606716 SPACESTORM and by the European Union's Horizon 2020 research and innovation programme under grant agreement No 637302 PROGRESS

**Special thanks to Dr. Elena Grigorenko (Space Research Institute, Moscow, Russia)  
for Cluster data analysis**

**Cluster 15th and Double Star 10th Anniversary  
Workshop 12---16 October 2015 Venice, Italy**



FINNISH  
METEOROLOGICAL  
INSTITUTE



# Plasma sheet electrons

The distributions of electrons with **1 to few hundreds of keVs** and their variations at **distances 6.6-11 Re** not sufficiently studied in detail.

**Modeling attempts:** *Jordanova and Miyoshi, 2005; Miyoshi et al., 2006; Chen et al., 2006; Jordanova et al., 2014.*

The electron flux at the keV energies is largely determined by convective and substorm-associated electric fields and varies significantly with geomagnetic activity driven by the solar wind [*Mauk and Meng, 1983; Kerns et al., 1994; Liemohn et al., 1998; Ganushkina et al., 2013, 2014*].

**Studies on keV electrons:** *Korth et al. [1999], Denton et al. [2005], Sicard-Piet et al. [2008], Denton et al. [2015]* (LANL MPA and SOPA electron data), *Friedel et al. [2001]* (Polar Hydra instrument), *Kurita et al. [2011]* (THEMIS spacecraft), *Asnes et al., [2008], Burin des Roziers et al. [2009]* (GEOTAIL and CLUSTER data)

In the near-Earth plasma sheet, continuous measurements of plasma sheet electrons are not available, in contrast to geostationary orbit

**No solar wind driven empirical relations** for electron fluxes or moments of electron distribution function which can be used easily for radiation belt modeling.

# Boundary conditions in the plasma sheet for modeling of keV electrons

Our previous studies [*Ganushkina et al.*, 2013, 2014]:

we set the model **boundary at  $10 R_E$**  and use the **kappa electron distribution** function.

Parameters of the kappa distribution function: **number density  $n$  and temperature  $T$**  in the plasma sheet given by the empirical model derived from Geotail data by TM03

*Tsyganenko and Mukai* [2003].

The **electron  $n$**  is assumed to be the same as that for ions in the TM03 model, but  **$T_e/T_i = 0.2$**  is taken into account (*Kaufmann et al.*, 2005; *Wang et al.*, 2012).

Applying this model for boundary conditions has a number of **limitations**:

- (1) Model was derived from Geotail data for ions (limited detector energy range  $<40\text{keV}$ ).
- (2) ratio  $T_e/T_i$  can vary during disturbed conditions.
- (3) at distances closer than  $10 R_E$ , the correlation between  $T_i$  and  $T_e$  might not exist at all and no certain ratio can be determined (*Runov et al.*, 2015).
- (4) simple  $\sin^2$  MLT dependence.

# Revision of boundary conditions in the plasma sheet using Cluster data

**Cluster data for ions and electrons used:**


**HIA**, from C1 and/or C3 (5 eV - 32 keV) and **PEACE** (0.7 eV - 26 keV).

For the Cluster magnetotail session in 2001-2005 we choose 70 orbits, which cross the CPS at  $-16\text{Re} < X_{\text{gsm}} < -5\text{Re}$  and  $|Y_{\text{gsm}}| < 11\text{Re}$ . The CPS was identified as a region where  $|B_x| \leq 5\text{nT}$ .

Totally from four Cluster satellites  
174558 moments (4s bins) of the CPS.

For each CPS moment the IMF and solar wind data ( $B_z$ ,  $N_{\text{sw}}$ ,  $V_{\text{sw}}$  in GSM) were determined by taking into account the time delay due to the solar wind propagation from ACE to Subsolar point.

Figure courtesy: E. Grigorenko







Electron density and temperature  
observed by Cluster and averaged  
for each (IMF Bz, Nsw) and  
(IMF Bz, Vsw) bin

Figure courtesy: E. Grigorenko

Ion density (assume Ni=Ne) and electron  
temperature  $T_e = T_i \cdot 0.26$  using  
Tsyganenko and Mukai (2003) model


**Tsyganenko - Mukai model**

### CLUSTER Observations



Measured electron temperature and number density different than those given by TM03 model

# Comparison of electron density from Cluster data and *Tsyganenko and Mukai [2003]* model



Tsyganenko and Mukai [2003] model does not describe well the electron density in the CPS.



Also lower correlation coefficients for ion number density from Cluster and from the model obtained than in original paper.



# Comparison of electron and ion temperatures from Cluster data and *Tsyganenko and Mukai* [2003] model

The ratio of  $T_i/T_e$  varies in the CPS in a wide range of values from 0.17 to 18.0 and the average  $\langle T_i/T_e \rangle = 3.8$ , thus  $T_e = T_i * 0.26$ .

Figure courtesy: E. Grigorenko



Reason of inconsistency is the selection criterion for the CPS intervals:

***Tsyganenko and Mukai* [2003] model include** only “quiet” CPS intervals without flapping, while we used all CPS intervals. Intervals of CPS with hot electrons ( $T_{e\_Cluster} > 2000$  eV) cannot be described.




# Revision of boundary conditions in the plasma sheet using THEMIS data

**THEMIS data for ions and electrons** used (2007-2014):


**ESA** (a few eV up to 25 (30) keV) and **SST** (25 keV- few eVs).

Then we computed the **plasma moments** using last calibration procedures. After synchronization of the solar wind data with THEMIS plasma moments we got ~66,000 datapoints at 1.5 min resolution.



# Comparison of ion temperatures from THEMIS data and *Tsyganenko and Mukai [2003]* model

Subset of the data with  $R=10-10.5 R_E$  used.



Correlation is very low ( $CC = 0.22$ ) in comparison with high correlation obtained for such comparison in TM03 paper ( $CC=0.7$ ).


Due to limited range of radial distances in our comparison while the whole range of the Geotail ( $R=10-50 R_E$ ) was used in TM03 model.

However, the points mostly fit the TM03 dependence confirming high quality of the THEMIS plasma data.

TM03 might systematically underestimate  $T_i$  for high-temperature part of the distribution due to limited energy range of Geotail spectrometer

# Comparison of number densities of electrons from THEMIS data and of ions from *Tsyganenko and Mukai* [2003] model

Subset of the data with  $R=10-10.5 R_E$  is used.




The correlation is 0.49  
(close to 0.56 by TM03) in spite of  
limited range of the radial distances.

**The TM03 equation for number  
density for ions can be used for  
electrons for IMPTAM simulation.**

However, we are working towards  
improving the model for Ne.

# Comparison of temperatures of electrons from THEMIS data and of ions from *Tsyganenko and Mukai* [2003] model

Subset of the data with  $R=10-10.5 R_E$  is used.



Red line:  $T_e = T_i$

Blue line:  $T_e = T_i/5$

If the relation  $T_e = T_i/5$  would have been valid in this region, the points would be distributed along blue line.

TM03 ion temperature shows **almost no correlation** with measured electron temperature.

Similar to *Runov* [2015] (private communication): there is **no correlation between  $T_i$  and  $T_e$  at geocentric distances closer than  $R=12R_E$** .

# Empirical model for electron temperature at 6-11 R<sub>e</sub> based on Cluster and THEMIS data (1)

It was found that **solar wind velocity** shows highest correlation with electron temperature for at 6-11 R<sub>E</sub>. IMF is of secondary importance.

Having experimented with time lag and duration of the average, we found that highest correlations are achieved for 1h delayed solar wind velocity.

$$V^* = V_{sw}(t-1h) / 400 \text{ km/s}$$


B<sub>s</sub> is averaged over dt= 1h 15 min and delayed 30min

$$BS = \left( \int_{t-1.75h}^{t-0.5h} BS * dt \right) / \Delta t \quad BS^* = BS / 2 \text{ nT}$$

B<sub>N</sub> is averaged over dt= 2h delayed 1h

$$BN = \left( \int_{t-3h}^{t-1h} Bn * dt \right) / \Delta t \quad BN^* = BN / 2 \text{ nT}$$

# Empirical model for electron temperature at 6-11 R<sub>e</sub> based on Cluster and THEMIS data




Model depends on coordinates  
 $R^* = R/10R_E$  and  $\Psi$

$$\Psi = \arctg\left(-\frac{Y_{GSM}}{X_{GSM}}\right)$$

The **correlation coefficient** for the whole  
range of distances  $R = 6-11 R_E$  is **0.61**.  
The mean absolute deviation is 2keV

$$T_e = \left[ \begin{aligned} &2.92 - 2.92 \cdot R^* + 0.538 \cdot \Psi - 0.707 \cdot R^* \cdot \Psi + 1.09 \cdot R^* \cdot \Psi^2 + \\ &(1.84 \cdot R^* + 1.35 \cdot \Psi^2 - 2.48 \cdot R^* \cdot \Psi^2) V_{SW}^* - 0.127 \cdot \Psi \cdot BS^* - 0.267 \cdot R^* \cdot BN^* \end{aligned} \right]^2$$

# Example for February 28 –March 3, 2013 storm





# Summary

1. keV electrons in the near-Earth geospace are of a key importance though not studied in details: as low energy electron seed population for radiation belts, as responsible for surface charging on satellites.
2. Source of low energy electrons is in the plasma sheet
3. Continuous measurements are not available, thus we need a model with time-dependent boundary conditions in the plasma sheet. TM03 developed for ions are not suitable for electrons.
4. Long-term data analysis from Cluster HIA (2001-2005) and THEMIS ESA and SST (2007-2014) for development of empirical relations for electron temperature and number density at distances 6-11 Re.
5. As a first version of the model:  
 $N_e = N_i$   
 $T_e$  is driven mainly by 1 hour delayed  $V_{sw}$  and  $B_s$ ,  $B_N$ .
6. Final version:  $N_e$  dependence and  $T_e$  with cold and hot populations